

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 August 2002 (08.08.2002)

PCT

(10) International Publication Number  
**WO 02/060945 A2**

(51) International Patent Classification<sup>7</sup>: **C07K 14/47**

(21) International Application Number: PCT/US01/49488

(22) International Filing Date:  
27 December 2001 (27.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/258,334 27 December 2000 (27.12.2000) US

(71) Applicant: **LEXICON GENETICS INCORPORATED**  
[US/US]; 4000 Research Forest Drive, The Woodlands, TX  
77381 (US).

(72) Inventors: **YU, Xuanchuan**; 7900 N. Stadium #101,  
Houston, TX 77030 (US). **MIRANDA, Maricar**; 1800 El  
Paseo, Apt. 1708, Houston, TX 77054 (US).

(74) Agents: **ISHIMOTO, Lance, K.** et al.; LEXICON GE-  
NETICS INCORPORATED, 4000 Research Forest Drive,  
The Woodlands, TX 77381 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,  
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA,  
ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

WO 02/060945 A2

(54) Title: NOVEL HUMAN ION CHANNEL PROTEIN AND POLYNUCLEOTIDES ENCODING THE SAME

(57) Abstract: Novel human polynucleotide and polypeptide sequences are disclosed that can be used in therapeutic, diagnostic, and pharmacogenomic applications.

NOVEL HUMAN ION CHANNEL PROTEIN AND  
POLYNUCLEOTIDES ENCODING THE SAME

The present application claims the benefit of U.S. Provisional Application Number 60/258,334 which was filed on December 27, 2000 and is herein incorporated by reference in its entirety.

1. INTRODUCTION

The present invention relates to the discovery, identification, and characterization of novel human polynucleotides encoding a protein that shares sequence similarity with mammalian cyclic nucleotide-gated ion channel proteins. The invention encompasses the described polynucleotides, host cell expression systems, the encoded proteins, fusion proteins, polypeptides and peptides, antibodies to the encoded proteins and peptides, and genetically engineered animals that either lack or overexpress the disclosed genes, antagonists and agonists of the proteins, and other compounds that modulate the expression or activity of the proteins encoded by the disclosed genes, which can be used for diagnosis, drug screening, clinical trial monitoring, the treatment of diseases and disorders, and cosmetic or nutraceutical applications.

2. BACKGROUND OF THE INVENTION

Ion channels are integral membrane proteins that mediate or facilitate the passage of ions across the lipid bilayer. Given that ion transport across the membrane often plays an important physiological role, ion channel proteins are proven drug targets.

### 3. SUMMARY OF THE INVENTION

The present invention relates to the discovery, identification, and characterization of nucleotides that encode a novel human protein, and the corresponding amino acid  
5 sequence of this protein. The novel human protein (NHP) described for the first time herein shares structural similarity with mammalian cyclic nucleotide-gated ion channels, and particularly CNG ion channel alpha subunits associated with sensory (olfactory and visual) signal transduction (see, for  
10 example, GENBANK Accession No. NM\_007724).

The novel human nucleic acid sequences described herein, encode a protein/open reading frame (ORF) of 664 amino acids in length (SEQ ID NO: 2).

The invention also encompasses agonists and antagonists  
15 of the described NHPs, including small molecules, large molecules, mutant NHPs, or portions thereof, that compete with native NHP, peptides, and antibodies, as well as nucleotide sequences that can be used to inhibit the expression of the described NHPs (e.g., antisense and ribozyme molecules, and  
20 open reading frame or regulatory sequence replacement constructs) or to enhance the expression of the described NHPs (e.g., expression constructs that place the described polynucleotide under the control of a strong promoter system), and transgenic animals that express a NHP sequence, or "knock-  
25 outs" (which can be conditional) that do not express a functional NHP. Knock-out mice can be produced in several ways, one of which involves the use of mouse embryonic stem cells ("ES cells") lines that contain gene trap mutations in a murine homolog of at least one of the described NHPs. When the unique  
30 NHP sequences described in SEQ ID NOS:1-2 are "knocked-out" they provide a method of identifying phenotypic expression of the particular gene as well as a method of assigning function to previously unknown genes. In addition, animals in which the

unique NHP sequences described in SEQ ID NOS:1-2 are "knocked-out" provide a unique source in which to elicit antibodies to homologous and orthologous proteins which would have been previously viewed by the immune system as "self" and therefore would have failed to elicit significant antibody responses.

Additionally, the unique NHP sequences described in SEQ ID NOS:1-2 are useful for the identification of protein coding sequence and mapping a unique gene to a particular chromosome. These sequences identify actual, biologically verified, and therefore relevant, exon splice junctions as opposed to those that may have been bioinformatically predicted from genomic sequence alone. The sequences of the present invention are also useful as additional DNA markers for restriction fragment length polymorphism (RFLP) analysis, and in forensic biology.

Further, the present invention also relates to processes for identifying compounds that modulate, *i.e.*, act as agonists or antagonists, of NHP expression and/or NHP activity that utilize purified preparations of the described NHPs and/or NHP product, or cells expressing the same. Such compounds can be used as therapeutic agents for the treatment of any of a wide variety of symptoms associated with biological disorders or imbalances.

#### 4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

The Sequence Listing provides the sequence of a NHP ORF encoding the described NHP amino acid sequence.

#### 5. DETAILED DESCRIPTION OF THE INVENTION

The NHP described for the first time herein is a novel protein whose transcript can be detected expressed in human testis cells.

The present invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, the expression products of such nucleotides, and: (a) nucleotides that encode mammalian homologs of the described genes, including the specifically described NHP, and the NHP products; (b) nucleotides that encode one or more portions of the NHP that correspond to functional domains, and the polypeptide products specified by such nucleotide sequences, including but not limited to the novel regions of any active domain(s); (c) isolated nucleotides that encode mutant versions, engineered or naturally occurring, of the described NHP in which all or a part of at least one domain is deleted or altered, and the polypeptide products specified by such nucleotide sequences, including but not limited to soluble proteins and peptides in which all or a portion of the signal sequence is deleted; (d) nucleotides that encode chimeric fusion proteins containing all or a portion of a coding region of NHP, or one of its domains (e.g., a receptor or ligand binding domain, accessory protein/self-association domain, etc.) fused to another peptide or polypeptide; or (e) therapeutic or diagnostic derivatives of the described polynucleotides such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy constructs comprising a sequence first disclosed in the Sequence Listing.

As discussed above, the present invention includes: (a) the human DNA sequences presented in the Sequence Listing (and vectors comprising the same) and additionally contemplates any nucleotide sequence encoding a contiguous NHP open reading frame (ORF) that hybridizes to a complement of a DNA sequence presented in the Sequence Listing under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M  $\text{NaHPO}_4$ , 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and

washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & Sons, Inc., NY, at p. 2.10.3) and encodes a functionally equivalent expression  
5 product.

Additionally contemplated are any nucleotide sequences that hybridize to the complement of a DNA sequence that encodes and expresses an amino acid sequence presented in the Sequence Listing under moderately stringent conditions, e.g., washing in  
10 0.2xSSC/0.1% SDS at 42°C (Ausubel et al., 1989, *supra*), yet still encodes a functionally equivalent NHP product. Functional equivalents of a NHP include naturally occurring NHPs present in other species and mutant NHPs whether naturally occurring or engineered (by site directed mutagenesis, gene  
15 shuffling, directed evolution as described in, for example, U.S. Patent No. 5,837,458). The invention also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequences.

Additionally contemplated are polynucleotides encoding a  
20 NHP ORF, or its functional equivalent, encoded by polynucleotide sequences that are about 99, 95, 90, or about 85 percent similar or identical to corresponding regions of the nucleotide sequences of the Sequence Listing (as measured by BLAST sequence comparison analysis using, for example, the GCG  
25 sequence analysis package using standard default settings).

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the described NHP gene (or coding region) nucleotide sequences. Such hybridization conditions may be  
30 highly stringent or less highly stringent, as described above. In instances where the nucleic acid molecules are deoxyoligonucleotides ("DNA oligos"), such molecules are

generally about 16 to about 100 bases long, or about 20 to about 80, or about 34 to about 45 bases long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence Listing. Such oligonucleotides can be used in conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc.

Alternatively, such NHP oligonucleotides can be used as hybridization probes for screening libraries, and assessing gene expression patterns (particularly using a micro array or high-throughput "chip" format). Additionally, a series of the described NHP oligonucleotide sequences, or the complements thereof, can be used to represent all or a portion of the described NHP sequences. An oligonucleotide or polynucleotide sequence first disclosed in at least a portion of one or more of the sequences of SEQ ID NOS: 1-2 can be used as a hybridization probe in conjunction with a solid support matrix/substrate (resins, beads, membranes, plastics, polymers, metal or metallized substrates, crystalline or polycrystalline substrates, etc.). Of particular note are spatially addressable arrays (i.e., gene chips, microtiter plates, etc.) of oligonucleotides and polynucleotides, or corresponding oligopeptides and polypeptides, wherein at least one of the biopolymers present on the spatially addressable array comprises an oligonucleotide or polynucleotide sequence first disclosed in at least one of the sequences of SEQ ID NOS: 1-2, or an amino acid sequence encoded thereby. Methods for attaching biopolymers to, or synthesizing biopolymers on, solid support matrices, and conducting binding studies thereon are disclosed in, *inter alia*, U.S. Patent Nos. 5,700,637, 5,556,752, 5,744,305, 4,631,211, 5,445,934, 5,252,743,

4,713,326, 5,424,186, and 4,689,405 the disclosures of which are herein incorporated by reference in their entirety.

Addressable arrays comprising sequences first disclosed in SEQ ID NOS:1-2 can be used to identify and characterize the temporal and tissue specific expression of a gene. These addressable arrays incorporate oligonucleotide sequences of sufficient length to confer the required specificity, yet be within the limitations of the production technology. The length of these probes is within a range of between about 8 to about 2000 nucleotides. Preferably the probes consist of 60 nucleotides and more preferably 25 nucleotides from the sequences first disclosed in SEQ ID NOS:1-2.

For example, a series of the described oligonucleotide sequences, or the complements thereof, can be used in chip format to represent all or a portion of the described sequences. The oligonucleotides, typically between about 16 to about 40 (or any whole number within the stated range) nucleotides in length can partially overlap each other and/or the sequence may be represented using oligonucleotides that do not overlap. Accordingly, the described polynucleotide sequences shall typically comprise at least about two or three distinct oligonucleotide sequences of at least about 8 nucleotides in length that are each first disclosed in the described Sequence Listing. Such oligonucleotide sequences can begin at any nucleotide present within a sequence in the Sequence Listing and proceed in either a sense (5'-to-3') orientation vis-a-vis the described sequence or in an antisense orientation.

Microarray-based analysis allows the discovery of broad patterns of genetic activity, providing new understanding of gene functions and generating novel and unexpected insight into transcriptional processes and biological mechanisms. The use of addressable arrays comprising sequences first disclosed in SEQ



ID NOS:1-2 provides detailed information about transcriptional changes involved in a specific pathway, potentially leading to the identification of novel components or gene functions that manifest themselves as novel phenotypes.

5        Probes consisting of sequences first disclosed in SEQ ID NOS:1-2 can also be used in the identification, selection and validation of novel molecular targets for drug discovery. The use of these unique sequences permits the direct confirmation of drug targets and recognition of drug dependent changes in  
10 gene expression that are modulated through pathways distinct from the drugs intended target. These unique sequences therefore also have utility in defining and monitoring both drug action and toxicity.

      As an example of utility, the sequences first disclosed  
15 in SEQ ID NOS:1-2 can be utilized in microarrays or other assay formats, to screen collections of genetic material from patients who have a particular medical condition. These investigations can also be carried out using the sequences first disclosed in SEQ ID NOS:1-2 *in silico* and by comparing  
20 previously collected genetic databases and the disclosed sequences using computer software known to those in the art.

      Thus the sequences first disclosed in SEQ ID NOS:1-2 can be used to identify mutations associated with a particular disease and also as a diagnostic or prognostic assay.

25        Although the presently described sequences have been specifically described using nucleotide sequence, it should be appreciated that each of the sequences can uniquely be described using any of a wide variety of additional structural attributes, or combinations thereof. For example, a given  
30 sequence can be described by the net composition of the nucleotides present within a given region of the sequence in conjunction with the presence of one or more specific oligonucleotide sequence(s) first disclosed in the SEQ ID NOS:

1-2. Alternatively, a restriction map specifying the relative positions of restriction endonuclease digestion sites, or various palindromic or other specific oligonucleotide sequences can be used to structurally describe a given sequence. Such  
5 restriction maps, which are typically generated by widely available computer programs (e.g., the University of Wisconsin GCG sequence analysis package, SEQUENCHER 3.0, Gene Codes Corp., Ann Arbor, MI, etc.), can optionally be used in conjunction with one or more discrete nucleotide sequence(s)  
10 present in the sequence that can be described by the relative position of the sequence relative to one or more additional sequence(s) or one or more restriction sites present in the disclosed sequence.

For oligonucleotide probes, highly stringent conditions  
15 may refer, e.g., to washing in 6x SSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as NHP gene antisense molecules, useful, for example, in NHP gene  
20 regulation and/or as antisense primers in amplification reactions of NHP gene nucleic acid sequences. With respect to NHP gene regulation, such techniques can be used to regulate biological functions. Further, such sequences may be used as part of ribozyme and/or triple helix sequences that are also  
25 useful for NHP gene regulation.

Inhibitory antisense or double stranded oligonucleotides can additionally comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil,  
30 hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil,

beta-D-galactosylqueosine, inosine, N6-isopentenyladenine,  
1-methylguanine, 1-methylinosine, 2,2-dimethylguanine,  
2-methyladenine, 2-methylguanine, 3-methylcytosine,  
5-methylcytosine, N6-adenine, 7-methylguanine,  
5 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil,  
beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil,  
5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-  
oxyacetic acid (v), wybutoxosine, pseudouracil, queosine,  
2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil,  
10 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid  
methylester, uracil-5-oxyacetic acid (v), 5-methyl-  
2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w,  
and 2,6-diaminopurine.

The antisense oligonucleotide can also comprise at least  
15 one modified sugar moiety selected from the group including but  
not limited to arabinose, 2-fluoroarabinose, xylulose, and  
hexose.

In yet another embodiment, the antisense oligonucleotide  
will comprise at least one modified phosphate backbone selected  
20 from the group including, but not limited to, a  
phosphorothioate, a phosphorodithioate, a phosphoramidothioate,  
a phosphoramidate, a phosphordiamidate, a methylphosphonate, an  
alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide  
25 is an  $\alpha$ -anomeric oligonucleotide. An  $\alpha$ -anomeric  
oligonucleotide forms specific double-stranded hybrids with  
complementary RNA in which, contrary to the usual  $\beta$ -units, the  
strands run parallel to each other (Gautier et al., 1987, Nucl.  
Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-  
30 methylribonucleotide (Inoue et al., 1987, Nucl. Acids Res.  
15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al.,  
1987, FEBS Lett. 215:327-330). Alternatively, double stranded

RNA can be used to disrupt the expression and function of a targeted NHP.

Oligonucleotides of the invention can be synthesized by standard methods known in the art, e.g. by use of an automated  
5 DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides can be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of  
10 controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. USA 85:7448-7451), etc.

Low stringency conditions are well-known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled  
15 sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual (and periodic updates thereof), Cold Spring Harbor Press, NY; and Ausubel et al., 1989, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley  
20 Interscience, NY.

Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using appropriately stringent conditions or by PCR. The identification and characterization of human genomic clones is helpful for  
25 identifying polymorphisms (including, but not limited to, nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single nucleotide polymorphisms), determining the genomic structure of a given locus/allele, and designing diagnostic tests. For example, sequences derived  
30 from regions adjacent to the intron/exon boundaries of the human gene can be used to design primers for use in amplification assays to detect mutations within the exons,

introns, splice sites (e.g., splice acceptor and/or donor sites), etc., that can be used in diagnostics and pharmacogenomics.

For example, the present sequences can be used in  
5 restriction fragment length polymorphism (RFLP) analysis to identify specific individuals. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification (as generally described in U.S.  
10 Patent No. 5,272,057, incorporated herein by reference). In addition, the sequences of the present invention can be used to provide polynucleotide reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for  
15 example, providing another "identification marker" (i.e., another DNA sequence that is unique to a particular individual). Actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments.

20 Further, a NHP gene homolog can be isolated from nucleic acid from an organism of interest by performing PCR using two degenerate or "wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be total  
25 RNA, mRNA, and/or cDNA obtained by reverse transcription of mRNA prepared from human or non-human cell lines or tissue known or suspected to express an allele of a NHP gene. The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequence of the desired NHP  
30 gene. The PCR fragment can then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment can be labeled and used to screen a cDNA

library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment can be used to isolate genomic clones via the screening of a genomic library.

PCR technology can also be used to isolate full length  
5 cDNA sequences. For example, RNA can be isolated, following  
standard procedures, from an appropriate cellular or tissue  
source (*i.e.*, one known, or suspected, to express a NHP gene).  
A reverse transcription (RT) reaction can be performed on the  
RNA using an oligonucleotide primer specific for the most 5'  
10 end of the amplified fragment for the priming of first strand  
synthesis. The resulting RNA/DNA hybrid may then be "tailed"  
using a standard terminal transferase reaction, the hybrid may  
be digested with RNase H, and second strand synthesis may then  
be primed with a complementary primer. Thus, cDNA sequences  
15 upstream of the amplified fragment can be isolated. For a  
review of cloning strategies that can be used, see *e.g.*,  
Sambrook *et al.*, 1989, *supra*.

A cDNA encoding a mutant NHP sequence can be isolated,  
for example, by using PCR. In this case, the first cDNA strand  
20 may be synthesized by hybridizing an oligo-dT oligonucleotide  
to mRNA isolated from tissue known or suspected to be expressed  
in an individual putatively carrying a mutant NHP allele, and  
by extending the new strand with reverse transcriptase. The  
second strand of the cDNA is then synthesized using an  
25 oligonucleotide that hybridizes specifically to the 5' end of  
the normal sequence. Using these two primers, the product is  
then amplified via PCR, optionally cloned into a suitable  
vector, and subjected to DNA sequence analysis through methods  
well-known to those of skill in the art. By comparing the DNA  
30 sequence of the mutant NHP allele to that of a corresponding  
normal NHP allele, the mutation(s) responsible for the loss or

alteration of function of the mutant NHP gene product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known to carry a mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, obesity, high blood pressure, connective tissue disorders, infertility, etc.), or a cDNA library can be constructed using RNA from a tissue known, or suspected, to express a mutant NHP allele. A normal NHP gene, or any suitable fragment thereof, can then be labeled and used as a probe to identify the corresponding mutant NHP allele in such libraries. Clones containing mutant NHP sequences can then be purified and subjected to sequence analysis according to methods well-known to those skilled in the art.

Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known, or suspected, to express a mutant NHP allele in an individual suspected of or known to carry such a mutant allele. In this manner, gene products made by the putatively mutant tissue can be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against a normal NHP product, as described below. For screening techniques, see, for example, Harlow, E. and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor, NY.

Additionally, screening can be accomplished by screening with labeled NHP fusion proteins, such as, for example, alkaline phosphatase-NHP or NHP-alkaline phosphatase fusion proteins. In cases where a NHP mutation results in an expression product with altered function (e.g., as a result of a missense or a frameshift mutation), polyclonal antibodies to NHP are likely to cross-react with a corresponding mutant NHP

expression product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well-known in the art.

The invention also encompasses (a) DNA vectors that  
5 contain any of the foregoing NHP coding sequences and/or their complements (*i.e.*, antisense); (b) DNA expression vectors that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences (for example, baculovirus as  
10 described in U.S. Patent No. 5,869,336 herein incorporated by reference); (c) genetically engineered host cells that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (d)  
15 genetically engineered host cells that express an endogenous NHP sequence under the control of an exogenously introduced regulatory element (*i.e.*, gene activation). As used herein, regulatory elements include, but are not limited to, inducible and non-inducible promoters, enhancers, operators and other  
20 elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include but are not limited to the cytomegalovirus (hCMV) immediate early gene, regulatable, viral elements (particularly retroviral LTR promoters), the early or late promoters of SV40 adenovirus, the  
25 *lac* system, the *trp* system, the *TAC* system, the *TRC* system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase (PGK), the promoters of acid phosphatase, and the promoters of the yeast  $\alpha$ -mating factors.

30 The present invention also encompasses antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists and agonists of a NHP, as well as compounds or



nucleotide constructs that inhibit expression of a NHP sequence (transcription factor inhibitors, antisense and ribozyme molecules, or open reading frame sequence or regulatory sequence replacement constructs), or promote the expression of a NHP (e.g., expression constructs in which NHP coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.).

The NHP or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, antibodies, antagonists and agonists can be useful for the detection of mutant NHPs or inappropriately expressed NHPs for the diagnosis of disease. The NHP proteins or peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for screening for drugs (or high throughput screening of combinatorial libraries) effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of NHP in the body. The use of engineered host cells and/or animals may offer an advantage in that such systems allow not only for the identification of compounds that bind to the endogenous receptor for an NHP, but can also identify compounds that trigger NHP-mediated activities or pathways.

Finally, the NHP products can be used as therapeutics. For example, soluble derivatives such as NHP peptides/domains corresponding to a NHP, NHP fusion protein products (especially NHP-Ig fusion proteins, i.e., fusions of a NHP, or a domain of a NHP, to an IgFc), NHP antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate or act on downstream targets in a NHP-mediated pathway) can be used to directly treat diseases or disorders. For instance, the administration of an

effective amount of soluble NHP, or a NHP-IgFc fusion protein or an anti-idiotypic antibody (or its Fab) that mimics the NHP could activate or effectively antagonize the endogenous NHP receptor. Nucleotide constructs encoding such NHP products can  
5 be used to genetically engineer host cells to express such products *in vivo*; these genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of a NHP, a NHP peptide, or a NHP fusion protein to the body. Nucleotide constructs encoding functional NHPs, mutant NHPs, as  
10 well as antisense and ribozyme molecules can also be used in "gene therapy" approaches for the modulation of NHP expression. Thus, the invention also encompasses pharmaceutical formulations and methods for treating biological disorders.

Various aspects of the invention are described in greater  
15 detail in the subsections below.

#### 5.1 THE NHP SEQUENCES

The cDNA sequence and the corresponding deduced amino acid sequence of the described NHP are presented in the  
20 Sequence Listing. The NHP nucleotides were obtained from clustered ESTs, genomic sequence (GENBANK Accession No. AF002992), and cDNAs made from human testis mRNA (Edge Biosystems, Gaithersburg, MD). The described NHP is apparently encoded on the human X chromosome.

25 The described novel human polynucleotide sequences can be used, among other things, in the molecular mutagenesis/evolution of proteins that are at least partially encoded by the described novel sequences using, for example, polynucleotide shuffling or related  
30 methodologies. Such approaches are described in U.S. Patent Nos. 5,830,721 and 5,837,458 which are herein incorporated by reference in their entirety.

NHP gene products can also be expressed in transgenic animals. Animals of any species, including, but not limited to, worms, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, birds, goats, and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate NHP transgenic animals.

Any technique known in the art may be used to introduce a NHP transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to pronuclear microinjection (Hoppe, P.C. and Wagner, T.E., 1989, U.S. Patent No. 4,873,191); retrovirus-mediated gene transfer into germ lines (Van der Putten et al., 1985, Proc. Natl. Acad. Sci., USA 82:6148-6152); gene targeting in embryonic stem cells (Thompson et al., 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer (Lavitrano et al., 1989, Cell 57:717-723); etc. For a review of such techniques, see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.

The present invention provides for transgenic animals that carry the NHP transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, i.e., mosaic animals or somatic cell transgenic animals. The transgene may be integrated as a single transgene or in concatamers, e.g., head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell-type by following, for example, the teaching of Lasko et al., 1992, Proc. Natl. Acad. Sci. USA 89:6232-6236. The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell-type of interest, and will be apparent to those of skill in the art.

When it is desired that a NHP transgene be integrated into the chromosomal site of the endogenous NHP gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous NHP gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous NHP gene (*i.e.*, "knockout" animals).

10 The transgene can also be selectively introduced into a particular cell-type, thus inactivating the endogenous NHP gene in only that cell-type, by following, for example, the teaching of Gu *et al.*, 1994, *Science*, 265:103-106. The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell-type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant NHP gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include but are not limited to Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and RT-PCR. Samples of NHP gene-expressing tissue, may also be evaluated immunocytochemically using antibodies specific for the NHP transgene product.

The present invention provides for "knockin" animals. Knockin animals are those in which a gene that the animal does not naturally have in its genome, is inserted. For example, when a human gene is used to replace its murine ortholog in the

mouse. Such knockin animals are useful for the *in vivo* study, testing and validation of, *intra alia*, human drug targets as well as for compounds that are directed at the same.

5                   5.2 THE NHP AND NHP POLYPEPTIDES

The described NHP, NHP polypeptides, peptide fragments, mutated, truncated, or deleted forms of the NHPs, and/or NHP fusion proteins can be prepared for a variety of uses. These uses include, but are not limited to, the generation of  
10 antibodies, as reagents in diagnostic assays, the identification of other cellular gene products related to NHP, as reagents in assays for screening for compounds that can be used as pharmaceutical reagents useful in the therapeutic  
15 treatment of mental, biological, or medical disorders and diseases. Given the similarity information and expression data, the described NHP can be targeted (by drugs, oligos, antibodies, etc, using drug screening assays similar to those described in, for example, U.S. Patent No. 6,048,850 herein  
20 treating diseases such as, for example, Alzheimer's disease, epilepsy, and Parkinson's disease.

The Sequence Listing discloses the amino acid sequence encoded by the described NHP ORF. The NHP displays an initiator methionine in a DNA sequence context consistent with  
25 a translation initiation site and several hydrophobic domains as typical of other membrane associated proteins.

The NHP amino acid sequence of the invention includes the amino acid sequence presented in the Sequence Listing as well as analogues and derivatives thereof. Further, corresponding  
30 NHP homologues from other species are encompassed by the invention. In fact, any NHP proteins encoded by the NHP nucleotide sequences described above are within the scope of the invention, as are any novel polynucleotide sequences

encoding all or any novel portion of an amino acid sequence presented in the Sequence Listing. The degenerate nature of the genetic code is well-known, and, accordingly, each amino acid presented in the Sequence Listing, is generically

5 representative of the well-known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such, as contemplated herein, the amino acid sequences presented in the Sequence Listing, when taken together with the genetic code (see, for example, Table 4-1 at page 109 of  
10 "Molecular Cell Biology", 1986, J. Darnell et al. eds., Scientific American Books, New York, NY, herein incorporated by reference) are generically representative of all the various permutations and combinations of nucleic acid sequences that can encode such amino acid sequences.

15 The invention also encompasses proteins that are functionally equivalent to the NHP encoded by the presently described nucleotide sequences as judged by any of a number of criteria, including, but not limited to, the ability to bind and cleave a substrate of a NHP, or the ability to effect an  
20 identical or complementary downstream pathway, or a change in cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation, etc.). Such functionally equivalent NHP proteins include, but are not limited to, additions or substitutions of amino acid residues within the amino acid  
25 sequence encoded by the NHP nucleotide sequences described above, but which result in a silent change, thus producing a functionally equivalent expression product. Amino acid substitutions can be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity,  
30 and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine,

tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged  
5 (acidic) amino acids include aspartic acid and glutamic acid.

A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the invention. Where, as in the present instance, the NHP peptide or polypeptide is thought to be membrane protein, the hydrophobic  
10 regions of the protein can be excised and the resulting soluble peptide or polypeptide can be recovered from the culture media. Such expression systems also encompass engineered host cells that express a NHP, or functional equivalent, *in situ*. Purification or enrichment of a NHP from such expression  
15 systems can be accomplished using appropriate detergents and lipid micelles and methods well-known to those skilled in the art. However, such engineered host cells themselves may be used in situations where it is important not only to retain the structural and functional characteristics of the NHP, but to  
20 assess biological activity, e.g., in certain drug screening assays.

The expression systems that may be used for purposes of the invention include, but are not limited to, microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with  
25 recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing NHP nucleotide sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing NHP nucleotide sequences; insect cell systems infected with recombinant virus expression  
30 vectors (e.g., baculovirus) containing NHP nucleotide sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV;

tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing NHP nucleotide sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs  
5 containing NHP nucleotide sequences and promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

In bacterial systems, a number of expression vectors may  
10 be advantageously selected depending upon the use intended for the NHP product being expressed. For example, when a large quantity of such a protein is to be produced for the generation of pharmaceutical compositions of or containing NHP, or for raising antibodies to a NHP, vectors that direct the expression  
15 of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which a NHP coding sequence may be ligated individually into the vector in frame with the *lacZ*  
20 coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors (Pharmacia or American Type Culture  
25 Collection) can also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The PGEX vectors are designed to include thrombin or factor Xa  
30 protease cleavage sites so that the cloned target expression product can be released from the GST moiety.



In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign polynucleotide sequences. The virus grows in *Spodoptera frugiperda* cells. A NHP coding sequence can be  
5 cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of NHP coding sequence will result in inactivation of the polyhedrin gene and production of non-  
10 occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted sequence is expressed (e.g., see Smith et al., 1983, J. Virol. 46: 584; Smith, U.S.  
15 Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the NHP nucleotide sequence of interest may be ligated to an adenovirus  
20 transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric sequence may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result  
25 in a recombinant virus that is viable and capable of expressing a NHP product in infected hosts (e.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted NHP nucleotide sequences. These  
30 signals include the ATG initiation codon and adjacent sequences. In cases where an entire NHP gene or cDNA, including its own initiation codon and adjacent sequences, is

inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (See Bitter et al., 1987, Methods in Enzymol. 153:516-544).

In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the expression product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and expression products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the expression product may be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, human cell lines.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell

lines which stably express the NHP sequences described above can be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media.

10 The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines

15 which express the NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the NHP product.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase

20 (Wigler et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska and Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy et al., 1980, Cell 22:817) genes, which can be employed in tk<sup>-</sup>, hgp<sup>-</sup> or ap<sup>-</sup> cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., 1980, Proc. Natl. Acad. Sci. USA 77:3567; O'Hare et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to

30 mycophenolic acid (Mulligan and Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., 1981, J. Mol.

Biol. 150:1); and hygro, which confers resistance to hygromycin (Santerre et al., 1984, Gene 30:147).

Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-8976). In this system, the sequence of interest is subcloned into a vaccinia recombination plasmid such that the sequence's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto  $\text{Ni}^{2+}$ -nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

Also encompassed by the present invention are fusion proteins that direct the NHP to a target organ and/or facilitate transport across the membrane into the cytosol. Conjugation of NHPs to antibody molecules or their Fab fragments could be used to target cells bearing a particular epitope. Attaching the appropriate signal sequence to the NHP would also transport the NHP to the desired location within the cell. Alternatively targeting of NHP or its nucleic acid sequence might be achieved using liposome or lipid complex based delivery systems. Such technologies are described in "Liposomes: A Practical Approach", New, R.R.C., ed., Oxford University Press, New York and in U.S. Patent Nos. 4,594,595, 5,459,127, 5,948,767 and 6,110,490 and their respective disclosures which are herein incorporated by reference in their entirety. Additionally embodied are novel protein constructs engineered in such a way that they facilitate transport of the

NHP to the target site or desired organ, where they cross the cell membrane and/or the nucleus where the NHP can exert its functional activity. This goal may be achieved by coupling of the NHP to a cytokine or other ligand that provides targeting  
5 specificity, and/or to a protein transducing domain (see generally U.S. applications Ser. No. 60/111,701 and 60/056,713, both of which are herein incorporated by reference, for examples of such transducing sequences) to facilitate passage across cellular membranes and can optionally be engineered to  
10 include nuclear localization..

Additionally contemplated are oligopeptides that are modeled on an amino acid sequence first described in the Sequence Listing. Such NHP oligopeptides are generally between about 10 to about 100 amino acids long, or between about 16 to  
15 about 80, or between about 20 to about 35 amino acids long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence Listing. Such NHP oligopeptides can be of any length disclosed within the above ranges and can initiate at  
20 any amino acid position represented in the Sequence Listing.

### 5.3 ANTIBODIES TO NHP PRODUCTS

Antibodies that specifically recognize one or more epitopes of a NHP, or epitopes of conserved variants of a NHP,  
25 or peptide fragments of a NHP are also encompassed by the invention. Such antibodies include but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')<sub>2</sub> fragments, fragments produced by a Fab expression  
30 library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

The antibodies of the invention may be used, for example, in the detection of NHP in a biological sample and may,

therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of NHP. Such antibodies may also be utilized in conjunction with, for example, compound screening schemes for the  
5 evaluation of the effect of test compounds on expression and/or activity of a NHP expression product. Additionally, such antibodies can be used in conjunction gene therapy to, for example, evaluate the normal and/or engineered NHP-expressing cells prior to their introduction into the patient. Such  
10 antibodies may additionally be used as a method for the inhibition of abnormal NHP activity. Thus, such antibodies may, therefore, be utilized as part of treatment methods.

For the production of antibodies, various host animals may be immunized by injection with the NHP, an NHP peptide  
15 (e.g., one corresponding to a functional domain of an NHP), truncated NHP polypeptides (NHP in which one or more domains have been deleted), functional equivalents of the NHP or mutated variant of the NHP. Such host animals may include but are not limited to pigs, rabbits, mice, goats, and rats, to  
20 name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including, but not limited to, Freund's adjuvant (complete and incomplete), mineral salts such as aluminum hydroxide or aluminum phosphate, chitosan, surface active substances such as  
25 lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Alternatively, the immune response could be enhanced by combination and or coupling with molecules such as keyhole  
30 limpet hemocyanin, tetanus toxoid, diphtheria toxoid, ovalbumin, cholera toxin or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, can be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated *in vitro* or *in vivo*. Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, Proc. Natl. Acad. Sci. USA, 81:6851-6855; Neuberger et al., 1984, Nature, 312:604-608; Takeda et al., 1985, Nature, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used (see U.S. Patent Nos. 6,075,181 and 5,877,397 which are herein incorporated by reference in their entirety). A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. Such technologies are described in U.S. Patent Nos. 6,075,181 and 5,877,397 and their respective disclosures which are herein incorporated by reference in their entirety. Also encompassed by the present

invention is the use of fully humanized monoclonal antibodies as described in US Patent No. 6,150,584 and respective disclosures which are herein incorporated by reference in their entirety.

5        Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778; Bird, 1988, Science 242:423-426; Huston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward et al., 1989, Nature 341:544-546) can be adapted to produce single chain antibodies against  
10 NHP expression products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments that recognize specific epitopes may  
15 be generated by known techniques. For example, such fragments include, but are not limited to: the F(ab')<sub>2</sub> fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab  
20 expression libraries may be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Antibodies to a NHP can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" a given NHP, using  
25 techniques well-known to those skilled in the art. (See, e.g., Greenspan & Bona, 1993, FASEB J 7(5):437-444; and Nissinoff, 1991, J. Immunol. 147(8):2429-2438). For example antibodies which bind to a NHP domain and competitively inhibit the binding of NHP to its cognate receptor can be used to generate  
30 anti-idiotypes that "mimic" the NHP and, therefore, bind and activate or neutralize a receptor. Such anti-idiotypic



antibodies or Fab fragments of such anti-idiotypes can be used in therapeutic regimens involving a NHP-mediated pathway.

Additionally given the high degree of relatedness of mammalian NHPs, the presently described knock-out mice (having  
5 never seen NHP, and thus never been tolerized to NHP) have a unique utility, as they can be advantageously applied to the generation of antibodies against the disclosed mammalian NHP (i.e., NHP will be immunogenic in NHP knock-out animals).

The present invention is not to be limited in scope by  
10 the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein  
15 will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims. All cited publications, patents, and patent applications are herein incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising at least 24 contiguous bases of nucleotide sequence first disclosed in the NHP sequence described in SEQ ID NO: 1.

2. An isolated nucleic acid molecule comprising a nucleotide sequence that:

- 10 (a) encodes the amino acid sequence shown in SEQ ID NO: 2; and
- (b) hybridizes under stringent conditions to the nucleotide sequence of SEQ ID NO: 1 or the complement thereof.

15 3. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes the amino acid sequence shown in SEQ ID NO: 2.

20 4. An isolated oligopeptide having a sequence of at least about 12 contiguous amino acids first disclosed in SEQ ID NO:2.

## SEQUENCE LISTING

&lt;110&gt; LEXICON GENETICS INCORPORATED

&lt;120&gt; Novel Human Ion Channel Protein and Polynucleotides Encoding the Same

&lt;130&gt; LEX-0291-PCT

&lt;150&gt; US 60/258,334

&lt;151&gt; 2000-12-27

&lt;160&gt; 2

&lt;170&gt; FastSEQ for Windows Version 4.0

&lt;210&gt; 1

&lt;211&gt; 1995

&lt;212&gt; DNA

&lt;213&gt; homo sapiens

&lt;400&gt; 1

atgaccgaaa	aaaccaatgg	tgtgaagagc	tccccagcca	ataatcacia	ccatcatgca	60
cctcctgcc	tcaaggccaa	tggcaaagat	gaccacagga	caagcagcag	gccacactct	120
gcagctgacg	atgacacctc	ctcagaactg	cagaggctgg	cagacgtgga	tgccccacag	180
caggggaagga	gtggcttccg	caggatagtt	cgcttggtgg	ggatcatcag	agaatgggcc	240
aacaagaatt	tccgagagga	ggaacctagg	cctgactcat	tcctcgagcg	tttctgtggg	300
cctgaactcc	agactgtgac	cacacaggag	ggggatggca	aaggcgacaa	ggatggcgag	360
gacaaaggca	ccaagaagaa	atgtgaacta	tttgtcttgg	acccagctgg	ggattggtac	420
tactgtctggc	tatttgtcat	tgccatggcc	gtcctttaca	actggtgcct	gctgggtggc	480
agagcctgct	tcagtgaact	acagaaaggc	tactacctgg	tgtggctggg	gctggattat	540
gtctcagatg	tgtgtctacat	tgcggacctc	ttcatccgat	tgcgcacagg	tttcctggag	600
caggggctgc	tgttcaaaga	taccaagaaa	ctgcgagaca	actacatcca	caccctgcag	660
ttcaagctgg	atgtggcttc	catcatcccc	actgacctga	tctattttgc	tgtggacatc	720
cacagccctg	aggtgcgctt	caaccgcctg	ctgcactttg	cccgcagtgt	tgagttcttt	780
gaccggacag	agacacgcac	caactaccct	aacatcttcc	gcatcagcaa	ccttgtcctc	840
tacatcttgg	tcatcatcca	ctggaatgcc	tgcactctatt	atgccatctc	caaattccata	900
ggctttgggg	tcgacacctg	ggttttacca	aacatcactg	accctgagta	tggctacctg	960
gctagggaat	acatctattg	cctttactgg	tccacactga	ctctcactac	cattggggag	1020
acaccacccc	ctgtaaagga	tgaggagtac	ctatttgtca	tctttgactt	cctgattggc	1080
gtcctcatct	ttgccaccat	cgtgggaaat	gtgggctcca	tgatctccaa	catgaatgcc	1140
acccgggcag	agttccaggc	taagatcgat	gccgtgaaac	actacatgca	gttccgaaag	1200
gtcagcaagg	ggatggaagc	caaggctcatt	agggtggttg	actacttggt	gaccaataag	1260
aagacagtgg	atgagcgaga	aattctcaag	aatctgccag	ccaagctcag	ggctgagata	1320
gccatcaatg	tccacttgtc	cacactcaag	aaagtgcgca	tcttccatga	ttgtgaggct	1380
ggcctgctgg	tagagctggg	actgaaactc	cgctctcagg	tcttcagtcc	tggggattac	1440
atgtgccgca	aaggggacat	cggcaaggag	atgtacatca	ttaaggaggg	caaactggca	1500
gtggtggctg	atgatgggtg	gactcagtat	gctctgctgt	cggctggaag	ctgctttggc	1560
gagatcagta	tccttaacat	taagggcagt	aaaatgggca	atcgacgcac	agctaataatc	1620
cgcagcctgg	gctactcaga	tctcttctgc	ttgtccaagg	atgatcttat	ggaagctgtg	1680
actgagtacc	ctgatgccaa	gaaagtccta	gaagagaggg	gtcgggagat	cctcatgaag	1740
gagggactgc	tggatgagaa	cgaagtggca	accagcatgg	aggtcgacgt	gcaggagaag	1800
ctagggcagc	tggagaccaa	catggaaacc	ttgtacactc	gctttggccg	cctgctggct	1860
gagtacacgg	gggcccagca	gaagctcaag	cagcgcacat	cagttcttga	aaccaagatg	1920
aaacagaaca	atgaagatga	ctacctgtct	gatgggatga	acagccctga	gctggctgct	1980
gctgacgagc	cataa					1995

<210> 2  
 <211> 664  
 <212> PRT  
 <213> homo sapiens

<400> 2

```

Met Thr Glu Lys Thr Asn Gly Val Lys Ser Ser Pro Ala Asn Asn His
 1           5           10           15
Asn His His Ala Pro Pro Ala Ile Lys Ala Asn Gly Lys Asp Asp His
      20           25           30
Arg Thr Ser Ser Arg Pro His Ser Ala Ala Asp Asp Asp Thr Ser Ser
      35           40           45
Glu Leu Gln Arg Leu Ala Asp Val Asp Ala Pro Gln Gln Gly Arg Ser
      50           55           60
Gly Phe Arg Arg Ile Val Arg Leu Val Gly Ile Ile Arg Glu Trp Ala
      65           70           75           80
Asn Lys Asn Phe Arg Glu Glu Glu Pro Arg Pro Asp Ser Phe Leu Glu
      85           90           95
Arg Phe Arg Gly Pro Glu Leu Gln Thr Val Thr Thr Gln Glu Gly Asp
      100           105           110
Gly Lys Gly Asp Lys Asp Gly Glu Asp Lys Gly Thr Lys Lys Lys Phe
      115           120           125
Glu Leu Phe Val Leu Asp Pro Ala Gly Asp Trp Tyr Tyr Cys Trp Leu
      130           135           140
Phe Val Ile Ala Met Pro Val Leu Tyr Asn Trp Cys Leu Leu Val Ala
      145           150           155           160
Arg Ala Cys Phe Ser Asp Leu Gln Lys Gly Tyr Tyr Leu Val Trp Leu
      165           170           175
Val Leu Asp Tyr Val Ser Asp Val Val Tyr Ile Ala Asp Leu Phe Ile
      180           185           190
Arg Leu Arg Thr Gly Phe Leu Glu Gln Gly Leu Leu Val Lys Asp Thr
      195           200           205
Lys Lys Leu Arg Asp Asn Tyr Ile His Thr Leu Gln Phe Lys Leu Asp
      210           215           220
Val Ala Ser Ile Ile Pro Thr Asp Leu Ile Tyr Phe Ala Val Asp Ile
      225           230           235           240
His Ser Pro Glu Val Arg Phe Asn Arg Leu Leu His Phe Ala Arg Met
      245           250           255
Phe Glu Phe Phe Asp Arg Thr Glu Thr Arg Thr Asn Tyr Pro Asn Ile
      260           265           270
Phe Arg Ile Ser Asn Leu Val Leu Tyr Ile Leu Val Ile Ile His Trp
      275           280           285
Asn Ala Cys Ile Tyr Tyr Ala Ile Ser Lys Ser Ile Gly Phe Gly Val
      290           295           300
Asp Thr Trp Val Tyr Pro Asn Ile Thr Asp Pro Glu Tyr Gly Tyr Leu
      305           310           315           320
Ala Arg Glu Tyr Ile Tyr Cys Leu Tyr Trp Ser Thr Leu Thr Leu Thr
      325           330           335
Thr Ile Gly Glu Thr Pro Pro Pro Val Lys Asp Glu Glu Tyr Leu Phe
      340           345           350
Val Ile Phe Asp Phe Leu Ile Gly Val Leu Ile Phe Ala Thr Ile Val
      355           360           365
Gly Asn Val Gly Ser Met Ile Ser Asn Met Asn Ala Thr Arg Ala Glu
      370           375           380
Phe Gln Ala Lys Ile Asp Ala Val Lys His Tyr Met Gln Phe Arg Lys
      385           390           395           400

```

Val Ser Lys Gly Met Glu Ala Lys Val Ile Arg Trp Phe Asp Tyr Leu  
 405 410 415  
 Trp Thr Asn Lys Lys Thr Val Asp Glu Arg Glu Ile Leu Lys Asn Leu  
 420 425 430  
 Pro Ala Lys Leu Arg Ala Glu Ile Ala Ile Asn Val His Leu Ser Thr  
 435 440 445  
 Leu Lys Lys Val Arg Ile Phe His Asp Cys Glu Ala Gly Leu Leu Val  
 450 455 460  
 Glu Leu Val Leu Lys Leu Arg Pro Gln Val Phe Ser Pro Gly Asp Tyr  
 465 470 475 480  
 Ile Cys Arg Lys Gly Asp Ile Gly Lys Glu Met Tyr Ile Ile Lys Glu  
 485 490 495  
 Gly Lys Leu Ala Val Val Ala Asp Asp Gly Val Thr Gln Tyr Ala Leu  
 500 505 510  
 Leu Ser Ala Gly Ser Cys Phe Gly Glu Ile Ser Ile Leu Asn Ile Lys  
 515 520 525  
 Gly Ser Lys Met Gly Asn Arg Arg Thr Ala Asn Ile Arg Ser Leu Gly  
 530 535 540  
 Tyr Ser Asp Leu Phe Cys Leu Ser Lys Asp Asp Leu Met Glu Ala Val  
 545 550 555 560  
 Thr Glu Tyr Pro Asp Ala Lys Lys Val Leu Glu Glu Arg Gly Arg Glu  
 565 570 575  
 Ile Leu Met Lys Glu Gly Leu Leu Asp Glu Asn Glu Val Ala Thr Ser  
 580 585 590  
 Met Glu Val Asp Val Gln Glu Lys Leu Gly Gln Leu Glu Thr Asn Met  
 595 600 605  
 Glu Thr Leu Tyr Thr Arg Phe Gly Arg Leu Leu Ala Glu Tyr Thr Gly  
 610 615 620  
 Ala Gln Gln Lys Leu Lys Gln Arg Ile Thr Val Leu Glu Thr Lys Met  
 625 630 635 640  
 Lys Gln Asn Asn Glu Asp Asp Tyr Leu Ser Asp Gly Met Asn Ser Pro  
 645 650 655  
 Glu Leu Ala Ala Ala Asp Glu Pro  
 660